


Listing of Claims

1-55 (Cancelled)

56. (Previously Presented) A method of inhibiting hyperproliferation of malignant or neoplastic cells, comprising treating the cells episodically with an antiproliferative amount of an active vitamin D compound which is a hypocalcemic vitamin D, with reduced risk of hypercalcemia; the cells expressing a vitamin D receptor, wherein the amount of active vitamin D is a high dose which is between about 10 μ g to about 200 μ g/dose given once per week to once every 12 weeks.

57. (Previously Presented) A method in accordance with claim 56, wherein the malignant cells are associated with cancers of the breast, colon, prostate, lung, pancreas, endometrium, liver, squamous cell carcinoma, myeloid leukemia, melanoma, retinoblastoma, sarcomas of the soft tissues or bone.

58. (Previously Presented) A method in accordance with claim 56, wherein the hypocalcemic vitamin D compound is a compound of formula (III):



wherein A¹ and A² each are hydrogen or together represent a carbon-carbon bond, thus forming a double bond between C-22 and C-23; R¹ and R² are identical or different and are

hydrogen, lower alkyl, lower fluoroalkyl, O-lower alkyl, lower alkenyl, lower fluoroalkenyl, O-lower alkenyl, O-lower acyl, O-aromatic acyl, lower cycloalkyl with the proviso that R^1 and R^2 cannot both be an alkenyl group, or taken together with the carbon to which they are bonded, form a C_3 - C_8 cyclocarbon ring; R^3 is lower alkyl, lower alkenyl, lower fluoroalkyl, lower fluoroalkenyl, O-lower alkyl, O-lower alkenyl, O-lower acyl, O-aromatic acyl or lower cycloalkyl; X^1 is hydrogen or hydroxyl, or, taken with R^3 , constitutes a bond when R^3 is an alkenyl group, and X^2 is hydrogen or hydroxyl, or, taken with R^1 or R^2 , constitutes a double bond.

59. (Previously Presented) A method in accordance with claim 56 wherein the active vitamin D is 1α -hydroxyvitamin D_2 or $1\alpha,24$ -dihydroxyvitamin D_2 .

60. (Previously Presented) A method in accordance with claim 56 wherein the active vitamin D is 1α -hydroxyvitamin D_4 ; $1\alpha,25$ -dihydroxyvitamin D_2 ; $1\alpha,24,25$ -trihydroxyvitamin D_2 ; $1\alpha,25$ -dihydroxyvitamin D_4 ; $1\alpha,24,25$ -trihydroxyvitamin D_4 ; 24 -hydroxyvitamin D_2 ; or 24 -hydroxyvitamin D_4 .

61. (Previously Presented) The method of claim 56 wherein the active vitamin D lacks a hydrocarbon moiety at the C-24 position.

62. (Previously Presented) A method in accordance with claim 61 wherein the active vitamin D is $1\alpha,25$ -dihydroxyvitamin D_3 or 1α -dihydroxyvitamin D_3 .

63. (Previously Presented) A method in accordance with claim 56 wherein the amount of the active vitamin D is administered to a human cancer patient, the amount of the active vitamin D effective to inhibit the hyperproliferation of the neoplastic cells.

64. (Previously Presented) The method of claim 63 wherein the amount of the vitamin D compound is administered parenterally or orally in combination with a pharmaceutically acceptable carrier.

65. (Previously Presented) A method in accordance with claim 64 wherein the amount of vitamin D compound is administered parenterally.

66. (Previously Presented) A method in accordance with claim 65 wherein the amount of vitamin D compound is administered intravenously.

67. (Previously Presented) A method of inhibiting hyperproliferation of malignant or neoplastic cells, comprising treating the cells by co-administering an antihyperproliferative amount of an active vitamin D compound and an effective amount of an agent which is an antineoplastic agent, a bone agent, an antihypercalcemic agent or combinations thereof, the cells expressing a vitamin D receptor, the antiproliferative amount of the active vitamin D compound being a dose between 10 μ g to about 200 μ g/dose administered on an episodic basis which is once per week to about once per 12 weeks.

68. (Previously Presented) A method in accordance with claim 67 wherein an amount of the active vitamin D compound and an amount of the agent are episodically co-administered to a human cancer patient, the amount of the active vitamin D effective to inhibit the hyperproliferation of the neoplastic cells.

69. (Previously Presented) A method in accordance with claim 67 wherein the agent is an antineoplastic agent.

70. (Previously Presented) A method in accordance with claim 69 wherein the antineoplastic agent is given episodically and the active vitamin D is given concurrently with the antineoplastic agent.

71. (Previously Presented) A method in accordance with claim 69 wherein the antineoplastic agent is an antimetabolite, an antimicrotubule agent, an alkylating agent, a platinum agent, an anthrocycline, a topoisomerase inhibitor, an antibiotic, any other antineoplastic agent or combinations thereof.

72. (Previously Presented) A method in accordance with claim 67 wherein the bone agent is a bisphosphonate.

73. (Previously Presented) A method in accordance with claim 67 wherein an active vitamin D compound, an antineoplastic agent and an antihypercalcemic agent are co-administered.